

12. 2344

III
7787

STIRPIUM RARIORUM MINUSQUE COGNITARUM

IN SICILIA

SPONTE PROVENIENTIUM

DESCRIPTIONES

NONNULLIS ICONIBUS AUCTAE

A U C T O R E

ANTONINO BIVONA BERNARDI

BARONE ALTAE TURRIS

ACCADEMIAE AECONOMICA FLORENTINAE SOC. CORR. etc, etc,

MANIPULUS I.



PANORMI
TYPIS REGII.

MDCCCXIII.



PD 9939

1. *Constitutio* *Imperialis* *de* *Corona* *Austriaca*

2. *Constitutio* *Imperialis* *de* *Corona* *Austriaca*

3. *Constitutio* *Imperialis* *de* *Corona* *Austriaca*

4. *Constitutio* *Imperialis* *de* *Corona* *Austriaca*

5. *Constitutio* *Imperialis* *de* *Corona* *Austriaca*

6. *Constitutio* *Imperialis* *de* *Corona* *Austriaca*

7. *Constitutio* *Imperialis* *de* *Corona* *Austriaca*

8. *Constitutio* *Imperialis* *de* *Corona* *Austriaca*

9. *Constitutio* *Imperialis* *de* *Corona* *Austriaca*

10. *Constitutio* *Imperialis* *de* *Corona* *Austriaca*

11. *Constitutio* *Imperialis* *de* *Corona* *Austriaca*

12. *Constitutio* *Imperialis* *de* *Corona* *Austriaca*

13. *Constitutio* *Imperialis* *de* *Corona* *Austriaca*

14. *Constitutio* *Imperialis* *de* *Corona* *Austriaca*

15. *Constitutio* *Imperialis* *de* *Corona* *Austriaca*


VIRO EXCELLENTISSIMO
JOSEPHO VINTIMILLIO
PULCHRI MONTIS DYNASTAE etc.

A REGIS CUBICULO
AB AUGUSTI UNIVERSI REGNI VICARIIS CONSILIIS
D. JANUARIY ORDINIS EQITI
MAGNOQUE HISPANARUM PROCERI.

ANTONINUS BIVONA

*Editurus in lucem harum stirpium descriptiones, tacitus
mecum ajebam: cui putem inscribendum quidquid hoc est
laboris ae studii mei? Viro certe omnium lectissimo, au-
ctoritate, splendore, dignitate praestanti; qui doctissimus
ipse doctrinas omnes, earumque cultores et amet et prove-
hat; quique Patriae caritate succensus, non ei tantum
commoda, quae a bonis artibus ac disciplinis proficiscun-
tur; sed bonis omnibus cumulatam felicitatem, posthabitatis
suis rationibus, comparare admittatur. Talem cum mihi ef-
fingerem virum; in Te incidisse magna cum animi mei
voluptate intellexi. Esto igitur Tu mihi Moeenas: tenuem
hunc ingenii mei fructum ea excipe humanitate, quae me
ad ubiores in dies proferendos possit excitare.*





AUCTORIS MONITUM.

Primi hujus Manipuli Praefationem alteri, qui propediem
in lucem veniet praeponendam reliqui. Liberius enim in
ea, quam in praesens licere arbitrer, rationes promam,
quibus opus a me incepturn nomine SIGULARUM PLANTARUM
CENTURIAE typis edere desierim, quin imo, ejus operis for-
matam prout nunc cernitur, immutare sim coactus. Hic
interim silentio praeterire haud possum, quos Jacobo E-
duardo Smith Linneanae Societatis Londinensis Praesidi
habeam grates, illo quo me ob rerum Botanicarum com-
merciū afficit honore, a quo maximam percipisse fateor
utilitatem, eamque quam plurimum in posterum auctun-
iri confido.



183. *On the first day of the month of April* *the King of Siam*
184. *sent his Ambassador to the King of Siam* *to demand*
185. *that he should give up his claim to the Kingdom of Siam*
186. *and that he should return to his native country*
187. *and that he should never return to Siam*
188. *and that he should never return to Siam*
189. *and that he should never return to Siam*
190. *and that he should never return to Siam*
191. *and that he should never return to Siam*
192. *and that he should never return to Siam*
193. *and that he should never return to Siam*
194. *and that he should never return to Siam*
195. *and that he should never return to Siam*
196. *and that he should never return to Siam*
197. *and that he should never return to Siam*
198. *and that he should never return to Siam*
199. *and that he should never return to Siam*
200. *and that he should never return to Siam*

CARDUUS ARGYROA.

*Carduus foliis decurrentibus pinnatifidis spinosis squarrosis,
calycibus aggregatis pedunculatis arachnoideo-villosis,
squamis rectis subulatis.*

*Radix annua, fusiformis, digitii medii crassitie, intus
alba, extus flavicans, fibrisque aucta numerosis.*

*Caulis a sesquipedali ad tres ferè pedes altus, erectus,
multiflorus, angulatus, foliosus, lanuginosus, alisque
spinosissimis, flexuosis, undique crispatus.*

*Folia plenè decurrentia, lanceolata, profundè pinnatifida
laciinis sinuato-dentatis, spinis validis flavescentibus
in apice dentium angulorumque armata, necnon spi-
nulis inaequalibus margine subciliata; supra laetè vi-
ridia, ambitu argenteo-glauco, vel (ex toto) albo-
nitore superfusa, subtus pallida, lanuginosa: radi-
calia et caulina inferiora petiolata, reliqua sessilia,
et quo magis angustata, eo orridius squarrosa.*

*Flores terminales axillaresque, erecti, bini, terni, vel
quaterni, pedunculati, pedunculis tomentosis, foliosis.*

*Calix subrotundus, Ayellanae ferè magnitudine, arach-
noideo-lanuginosus, squamis imbricatus erectis, su-
bulatis, spina molli apice terminatis.*

*Corollæ quinquefidae, calyce longiores, dilutè purpu-
reæ, disco saepè albae.*

Antherae in intimis flosculorum seriebus semper polliniferae, coeruleae, in reliquis interdum steriles, albidae.
Stylus et Stigmata purpurea.

Semina compressa, pappo instructa piloso, albicante, scabro, demum deciduo.

Receptaculum pilosum.

Panormi in aggeribus et ad vias.

Annua. Floret *Aprili*, *Majo*.

VICIA LEUCANTHA.

Vicia pedunculis multifloris folio brevioribus; foliolis ellipticis villosis, stipulis semisagittatis linearis-subulatis dentatis.

Tota planta villosa.

Radix tenuis.

Caules debiles, lineam crassi, sesquipedem longi, tetragonoi, sulcati, cirrhis scandentes trifidis vel quadrifidis.

Foliola 14-20, elliptica cum mucronulo nigrescente, opposita vel alterna, 6-lineas longa, 2 $\frac{1}{2}$ -lineas lata, superiora sensim magnitudine decrescentia.

Stipulae semisagittatae, angustissimae, linearis-subulatae, dentatae, dentibus acutis.

Pedunculi folio breviores, 2-9flori, floribus secundis parvis.

Calix corollà brevior, campanulatus, villosus, basi purpurascens, quinquedentatus, dentibus setaceis subaequalibus.

Vexillum glabrum, emarginatum, albo-carneum, intus lineis coeruleis striatum.

Alae albae, obtusae, glabrae, vexillo paulo breviores.

Carina alis duplo brevior, apice nigra.

Stylus inflexus, versus apicem supra et subtus villosus.

Stigma capitatum.

*Legumina pendula, compressa, venosa, 14-lineas longa,
5 $\frac{1}{2}$ · lineas lata, villis tecta adpressis, rariusculis.*

Semina quinque.

Habitat circa Panormum in pascuis.

Annua. Floret Martio, Aprili.

ORCHIS BRANCIFORTII.

Tab. I. Fig. I.

Orchis labello trilobo plano, lobis integerrimis inaequalibus, cornu filiformi germen subaequante, petalis exterioribus patentissimis.

Orchis parva maculata purpureo flore Culicis effigie Bon.

t. 40 - Cup. Panph. 2 t. 118.

Radix testiculata, tuberibus ovatis, parvis.

Caulis 4-6 pollicaris, erectus, subangulatus, infernè foliosus, supernè nudus purpurascens.

Folia radicalia pauca, patentia, laetè viridia, (aliquando maculata) oblonga vel lanceolata, sesquipolllicem vel duos pollices longa, semipolllicem ferè lata; caulina erecta vaginantia margine purpurascens.

Spica laxa, pyramidata floribus parvis, roseis, glabermissis, inodoris.

Petala tria exteriora in triangulum patentia, elliptica, subaequalia, enervia.

Opercula petalis duplo ferè breviora, concava, ovata, in cucullum conniventia.

Labellum petalis paulo brevius, trilobum, planum, binis punctis purpureis ad basin plerumque notatum.

Lobi lineares, inaequales, patentes, integerrimi: laterales breviores, angustissimè lineares, apice subtruncati, medius latior, apice in angulum acutiusculum producto.

Columna antherifera brevissima, purpurea, mutica.

Nectarii cornu germine paulo brevius, filiforme subincurvum.

Bracteae purpureae, lanceolatae, germen vaginantes lineare angulatum duplo longius.

Habitat in montibus circa *Panormum*.

Perennis. Floret Martio, Aprili.

AERGULI BRANCIFORTE

BUTERAE PRINCIPI

Viro Nobilissimo atque Generosissimo, Botanices Siculae Fautori ac Patrono meo Colendissimo in grati animum et summae observantiae perpetuum documentum hanc novam plantam inscribere, ejusque perillustri cognomine exornare mihi fas esse duxi. Quam veniam utilium mili dent exteri Botanici; Siculique cives sibi sumant exemplum, atque hunc veluti gloriolae cuiusdam igniculum alere studeant atque in dies augescere.

ANTHOXANTHUM GRACILE.

Tab. I. Fig. II.

Anthoxanthum spica brevi pauciflorâ, flosculis aristâ duplo brevioribus, culmo filiformi.

Radix fibrosa, parva, annua, odorata odore Anthox. odorati.

Culmus 3-5 pollicaris, filiformis, erectus, supernè nudus scabriusculus tenuissimè striatus, infernè foliosus, ramosus, bi- vel trinodis.

Folia brevia, plana, ciliata, paginâ superiori villosa, inferiori striata, glabra; supremum vaginâ brevius. Vaginae parum ventricosae, subangulatae, plerumque glabrae, quandoque retrorsum villosae, ore nudæ. Stipula retusa, brevis, lacera vel denticulata.

Spica Aveneformis, terminalis, solitaria, erecta, brevis, pauciflora, rachi flexuosa.

Flores lanceolati, quatuor lineas (exclusâ aristâ) longi, per paria alterni, altero pedicellato, pedicello pileo.

Calix e glumis duabus acutis, quandoque aliquanto pilosis, margine membranaceis, apice scabris: exteriori uninervi duplo breviori, interiori trinervi, nervo medio in aristam brevem producto.

Corolla biglumis , setosa , ferruginea , calyce brevior .

Glumae subaequales acutae , aristatae , aristâ unâ dorsali , calyce breviori , aliâ basilarî geniculatâ , calyce duplo longiori , infernè glabrâ fuscâ , supernè scabrâ , viridi .

Nectarium bivalve german et filamentorum basin vaginans.

Stamina duo exerta .

Stigmata pariter exerta filiformia , villosa .

Habitat in montibus circa *Panormum* .

Annua . Floret *Aprilis* .

OROBANCHE FOETIDA.

Orobanche caule simplicissimo, corollâ inflatâ, labii inferioris lacinii rotundatis inaequalibus, argutè dentatis, staminibus basi intus villosis, stylo glanduloso.

Orobanche foetida Poiret Voy. 2 p. 196 - Desf. Atl. 2
p. 59 t. 144.

Orobanche major β Vahl. Symb. 2 p. 70.

β Lutea.

Tota planta pilis tecta capitatis, a pede ad sesquipedem alta, colore saepius atro-purpureo, aliquando flavo.

Radix parasitica, subbulbosa, solida, pollicem in diametro lata, extus squamosa.

Caulis simplicissimus, subflexuosus, angulatus, squamis vestitus sparsis, aridis, fuscis, lanceolatis, acuminatis.

Flores numerosi 6-8 lineas longi in spica terminali simplici.

Bracteae solitariae, florae longiores, calyci adpressae, acuminę reflexo.

Calycis foliola duo, distincta, oblonga, corollâ breviora, semibifida, lacinii inaequalibus, setaceo-acuminatis.

Corolla bilabiata, ventricosa, argutè dentata, dorso carinata. Labium superius fornicateum, bilobum, lobis rotundatis margine reflexis, inferius trilobum, lobis inaequalibus, distantibus, rotundis, concavis vel

planis , apice aliquando appendiculatis , sive denticulo productiori instructis . Palatum lucidum , sulcatum .

Stamina inclusa , fundo corollae inserta , superne pubescentia , inferne glabra , basi intus valde villosa . Antherae glabrae , bilobae , lobis spinulâ terminatis .

Stylus purpurascens , pubescens , staminibus longior , incurvus .

Stigma dilatatum , bilobum , flavum , subinde exertum .

Germen oblongum , basi glandulosum , apice pubescens .

Habitat in montibus circa *Panormum* ad radices *Spartii* .

Perennis . Floret *Majo* .

BARKAUSIA HYEMALIS.

T. II.

*Barkausia foliis radicalibus runcinatis, dentatis, caulinis
amplexicaulibus hastato-saggittatis, basi pinnatifidis,
calycibus hirtis, squamis exterioribus ovato-lanceolatis.*

*Hieracium hibernum Cichorii folio tenui Cup. Hort. Cath.
p. 95.*

*Hieracium hiemale Cichorii folio hirtum Cup. Panph. t.
t. 12 et 171 - Bonan. t. 125.*

Radix fusiformis, ramosa, extus nigrescens.

*Caules modo erecti, modo basi decumbentes, a semipede
ad sesquipedem alti, angulati, sulcati, ramosi, pur-
purascentes, pilis hirsuti simplicibus nigricantibus.*

*Folia glabra vel leviter hirsuta: radicalia petiolata, non-
nunquam integra, saepius pinnatifida, runcinata, lo-
bis acutis, inaequaliter dentatis, extremo majori
quandoque obtuso, caulina amplexicaulia, hastato-
sagittata, pinnatifido-dentata.*

*Flores corymboso-paniculati, citrini, semiflosculis exterio-
ribus dorso purpurascentibus.*

*Calyx tomentosus foliolis carinatis, pilis nigris hirsutis,
exterioribus ovato-lanceolatis, margine membranaceis,
subinde laxis.*

Semina flava , sulcata , gracilia , sub lente scabra , pap.
po simplici , stipitato , glabro .

Receptaculum areolatum , areolis margine villosiusculis ,
medio mammillatis .

Habitat circa *Panormum* in pratis et pascuis , muris te-
ctisve .

Perennis . Floret *Hyeme* et *Vere* .

T.I.



